



TEMPO[®] SC

ULTRA

Insecticide

For Pest Management Professionals and Commercial Use Only.

ACTIVE INGREDIENT: β - Cyfluthrin,
Cyano(4-fluoro-3-phenoxyphenyl)methyl
3-(2,2-dichloroethenyl)-2,2-dimethyl-
cyclopropanecarboxylate 11.8%
OTHER INGREDIENTS..... 88.2%
100.0%

Contains 1 lb cyano(4-fluoro-3-phenoxyphenyl)methyl 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate per gallon.

SHAKE WELL BEFORE USING.

EPA Reg. No. 432-1363

STOP - Read the label before use.
Keep out of reach of children.

CAUTION

Net Contents: 240 mL (8.12 fl oz)

4031958

3549856C 100504AV1

BACKED
by **BAYER**..

FIRST AID

If on skin:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
If in eyes:	• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. The Bayer Environmental Science Emergency Response Telephone No. is 1-800-334-7577.

Note To Physician: No specific antidote is available. Treat the patient symptomatically.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Remove from premises or tightly cover fish tanks and disconnect aerators when applying indoors where such containers are present. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Apply this product only as specified on this label.

This pesticide is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply TEMPO SC ULTRA Insecticide or allow it to drift onto crops or weeds on which bees are actively foraging. Additional information may be obtained by consulting your Cooperative Extension Service.

PRO FACTS	
• Quick Knockdown & Residual Control • Low Odor • Non-staining • No phytotoxicity on a wide range of ornamentals	
TEMPO SC ULTRA	• Suspension concentrate containing 1 lb β-cyfluthrin per gallon
CONTROLS	• Indoor Pests - General Household Pests - Pantry and Stored Product Pests - Food Processing Pests • Outdoor/ Perimeter Pests • Wood-Infesting Pests • Turf and Ornamental Pests For list of specific insects, see inside.

(continued)

WHERE TO APPLY	Apply where pests have been seen or found, or can find shelter. • In and around buildings and structures • On residential, commercial and recreational areas of turf • On ornamentals in landscapes and interior plantscapes
COVERAGE	• Makes up to 30 gals of finished spray solution for indoors. • Treats up to 30,000 sq. ft. outdoors.
MIXING INFORMATION	• Mix the appropriate amount of TEMPO SC ULTRA with water. • May tank mix with other pesticides currently registered for similar uses, unless specifically prohibited.
HOW TO APPLY	• General surface spot, mist, or crack & crevice application. • Broadcast application to turf and foliar application to ornamentals.
RE-APPLY	• Every 7 to 10 days, if needed. • For turf, re-apply if needed, but not more than 6 times per year.
RE-ENTRY 	• Avoid contact with treated surfaces until dry. • People and pets may re-enter after the treated area is dry.

(continued)

APPLICATION RESTRICTIONS	Do not water the treated area to the point of run-off. Do not make applications during rain. All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses: (1) Treatment to soil or vegetation around structures; (2) Applications to lawns, turf, and other vegetation; (3) Applications to building foundations, up to a maximum height of 3 feet. Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications, only. Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.
QUESTIONS 	For questions or comments, call toll-free 1-800-842-8020.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not formulate this product into other end-use products.

STRUCTURAL PESTS

FOR RESIDUAL PEST CONTROL IN AND AROUND BUILDINGS, STRUCTURES, FOOD/FEED HANDLING ESTABLISHMENTS AND MODES OF TRANSPORT

Use TEMPO SC ULTRA as a general surface, spot, mist, or crack and crevice treatment in and around buildings and structures. This includes schools, factories, stores, warehouses, hospitals, food processing facilities, and modes of transport. For a list of common facilities where this product may be used, see Appendix.

INDOOR PESTS

GENERAL HOUSEHOLD PESTS

<u>Crawling Pests</u>	Clothes moths	Earwigs	Sowbugs	Hornets
Ants (except Pharaoh)	Cluster flies	Firebrats	Spiders	Midges
Bedbugs	Cockroaches	Millipedes	<u>Flying Pests</u>	Moths
Carpet beetles	Crickets	Pillbugs	Flies	Wasps
Centipedes	Darkling beetles	Silverfish	Gnats	Yellow jackets

PANTRY & STORED PRODUCT PESTS

<u>Beetles</u>	Dermestid	Lesser grain borer	Sawtoothed	Indian meal
(exposed adults & immature stages)	Drugstore	Lesser mealworm	grain	Mediterranean
Cadelle	Granary weevil	Merchant grain	Warehouse	flour
Cigarette	Hide	Mealworm	<u>Moths</u>	
Confused flour	Larder	Red flour		
	Leather	Rice weevil		

(continued)

FOOD PROCESSING PESTS

Fruit flies
Phorid flies
Sciariid flies

Also includes:
General Household Pests
Pantry & Stored Product Pests

WHERE TO APPLY

Apply where pests have been seen or found, or can find shelter. Use a general surface, spot, mist, or crack and crevice application.

USE RATE

0.025% -- 8 ml (0.27 fl oz) per gal water
0.05% -- 16 ml (0.54 fl oz) per gal water

- Use high rate for severe infestations, faster knockdown, and longer residual.
- Always use the high rate for bees, wasps, yellow jackets and hornets.

HOW TO APPLY (Also see APPLICA- TION FOR INDOOR PESTS)

- Remove or tightly cover fish tanks and disconnect aerators during application.
- Remove pets during application and until treated surfaces are completely dry.
- Avoid contact with treated surfaces until dry.
- Wear safety glasses, goggles, or face shield and a dust/mist respirator when treating overhead areas.

General Surface Applications

- Use a low-pressure system (do not exceed 50 psi at nozzle tip), with a fan-type nozzle to apply the dilution uniformly.
- In the home, all food processing surfaces and utensils should be covered or thoroughly washed following treatment. Cover exposed food or remove from area being treated.

Spot, Mist, or Crack and Crevice Applications

- Apply directly into voids, cracks and crevices using spray or foam equipment.
- Spray: Use a low-pressure system (do not exceed 50 psi at nozzle tip), with a pinpoint or variable pattern nozzle or applicator equipment that delivers low volume treatments, such as the Micro-Injector® or Actisol®.

(continued)

	• Foam: Use appropriate foam generating equipment to supply a sufficient volume of foam. Mix the dilution with manufacturer's recommended volume of foaming agent.
WHEN TO APPLY	• When pests are seen or found • Re-apply every 7 to 10-days if needed.
MIXING INFORMATION	• Add product when filling spray tank with water; shake or agitate mixture. • Diluted spray mixture can be stored overnight. Agitate before using. • Can be tank mixed with other pesticides currently registered for similar uses. To be sure of physical compatibility, first prepare a sample in a small jar using the proper proportions of water and products. • Follow the most restrictive of labeling directions and precautions. Do not exceed label rates.
RESTRICTIONS	• Do not treat entire area of floor or floor coverings. • Do not apply where electrical short circuits can occur. Use a dust or dry bait in these areas. • Do not apply in occupied hospital and/or nursing home rooms. Occupants may return when surfaces have dried. • Do not apply in occupied classrooms. • Do not apply to furniture or upholstery where prolonged contact by humans will occur. • Do not apply as a space spray.

ADDITIONAL DIRECTIONS FOR FOOD AND FEED HANDLING ESTABLISHMENTS

WHERE TO APPLY	Applications may be made to both food/feed and non-food areas of food and feed handling establishments. Food/Feed handling establishments are defined as places other than private residences in which exposed food/feed is held, processed, prepared, or served. Includes areas for: • receiving, storing and packing (canning, bottling, wrapping, boxing) food • preparing food • edible waste storage • enclosed processing systems (mills, dairies, edible oils, syrups) of food. Serving areas where food is exposed and the facility is in operation are also considered food areas.
USE RATE	• Apply as a general surface spot, mist, or crack & crevice application. 0.025% -- 8 ml (0.27 fl oz) per gal water 0.05% -- 16 ml (0.54 fl oz) per gal water • Use high rate for severe infestations, faster knockdown, and longer residual.
HOW TO APPLY (Also see INDOOR PESTS)	<u>General Surface Applications</u> Cover or remove all food processing and handling equipment before treatment. After applying where food is commercially prepared or processed, wash all handling and processing equipment, benches, shelving, and other surfaces which food will contact. Rinse thoroughly with clean, fresh water. Do not apply to surfaces in food-handling areas when facility is in use or when foods are exposed. Do not apply directly to food. <u>Spot or Crack & Crevice Applications</u> Spot or crack and crevice application is permitted while facility is in use. Cover or remove food from area being treated. Do not apply directly to food or food handling surfaces.

APPLICATION FOR INDOOR PESTS	
GENERAL HOUSEHOLD PESTS Cockroaches, Crickets, Silverfish, Spiders, and others	• Use a coarse, low-pressure spray. • Treat entry points such as around water pipes, doors, windows, and eaves. Treat areas where pests normally feed or hide such as baseboards, corners, around water pipes, under refrigerators, cabinets, sinks, and stoves, being sure to spray cracks & crevices. Spot treat floor or rugs beneath furniture, in closets, and storage areas, but do not apply to entire floor area. • Do not use as a “space spray”. • IGR Note: This product can be tank mixed with an insect growth regulator (IGR) for extended cockroach control.
ANTS	• Apply to ant trails on walls, tile, baseboards, around pipes, under cabinets, and other places where ants are active. • Treat entry points such as windows and doors and the outdoor perimeter. • For best results, treat nests when possible.
PANTRY AND STORED PRODUCT PESTS	• For best results, clean area before treating. Remove and destroy infested food. Discard food accidentally contaminated with spray. • For cupboards, shelving, or food storage areas: Remove utensils, exposed food, shelf paper, and other items before spraying. When dried, cover shelves with clean paper before replacing items.
COMMERCIAL STORED PRODUCT PESTS	• Commercial storage or transportation: Clean and treat warehouses, production facilities, storage areas, rail cars, truck beds, and other areas where products are stored before filling with the product.
PESTS IN AIRCRAFT	• Do not use in aircraft cabins. • May be applied to carpet or upholstery in the cargo area only.

(continued)

PESTS IN LIVESTOCK HOUSING STRUCTURES including poultry houses & pet kennels	• Apply as a general surface and/or a crack and crevice spray. • For best results, clean facilities before applying and treat outdoor perimeter areas. • Do not apply indoors where animals other than cattle or horses are present. • Do not apply to animal feedstuffs or watering equipment.
FLYING INSECTS	• Apply to surfaces where pests collect or rest.

STRUCTURAL PESTS

OUTDOOR/PERIMETER PESTS

Ants	Cluster flies	Flies	Mosquitoes	Spiders
Bees	Crickets	Gnats	Moths	Ticks
Boxelder bugs	Elm leaf beetles	Ground beetles	Pillbugs	Wasps
Carpenter ants	Fire ants	Hornets	Scorpions	Yellow jackets
Centipedes	Firebrats	Midges	Silverfish	
Cockroaches	Fleas	Millipedes	Sowbugs	

WHERE TO APPLY	• Apply where pests have been seen or found, or can find shelter. • Apply to buildings, porches, patios, garages, fences and other areas where pests are found. • To prevent infestation, treat the building foundation, walls, and around doors, windows, and soffits where pests are active or may enter or hide; also treat the soil, turf, or other substrates adjacent to buildings.
-----------------------	---

(continued)

USE RATE	• 8 to 16 ml (0.27 to 0.54 fl oz) per 1000 sq ft - See Table 1 for mixing directions. • For fire ants and flying pests: 0.025% -- 8 ml (0.27 fl oz) per gal water 0.05% -- 16 ml (0.54 fl oz) per gal water • Always use the high rate for fire ants, bees, wasps, yellow jackets and hornets. • Use high rate for severe infestations, faster knockdown, and longer residual.
HOW TO APPLY (Also see APPLICATIONS FOR OUTDOOR/ PERIMETER PESTS)	<u>Perimeter Treatment:</u> To help prevent infestation of structures, treat area adjacent buildings in a band up to 10 feet wide. Also treat the building foundation to a height of 2 to 3 feet in areas where pests are active or can find entrance. • Apply just enough dilution to adequately cover the area without excessive dripping or runoff. Volume can vary depending on the surface type treated. For example, mulch requires more spray than bare soil. If mulch is heavy, rake it back to expose soil before treating. • Avoid contact with treated surfaces until dry.
WHEN TO APPLY	• When pests are seen or found • Re-apply every 7 to 10 days if needed.
MIXING INFORMATION	• Add product when filling spray tank with water; shake or agitate mixture. • Diluted spray mixture can be stored overnight. Agitate before using. • Can be tank mixed with other pesticides currently registered for similar uses. To be sure of physical compatibility, first prepare a sample in a small jar using the proper proportions of water and products. • Follow the most restrictive of labeling directions and precautions. Do not exceed label rates.

Table 1: Mixing Directions for Outdoor/Perimeter Applications

Application Volume: gals/1,000 sq ft	Use Rate: Milliliters/ 1,000 sq ft	Milliliters of TEMPO SC ULTRA Diluted to These Volumes of Finished Spray			
		3 gals	10 gals	15 gals	25 gals
2	8	12	40	60	100
	16	24	80	120	200
5	8	--	16	24	40
	16	--	32	48	80
10	8	--	8	12	20
	16	--	16	24	40

Formula for Determining the Percent Concentration in the Finished Spray

$$\frac{11.8 \times (\text{Milliliters of TEMPO SC ULTRA})}{(\text{gals of finished spray}) \times 3,785} = \text{Percent Concentration}$$

APPLICATION FOR OUTDOOR/PERIMETER PESTS**ANTS**

Locate and treat ant nests, if possible. Follow trails of foraging workers, check under objects near the structure, and inspect other likely nest sites. If ants are nesting in the soil, thoroughly drench or inject the nest and surrounding soil. Treat ant trails around windows and doors, along sidewalks, or other areas where seen. Apply a perimeter treatment as described above.

(continued)

BEEES, WASPS, YELLOW JACKETS, HORNETS	Use Rate: 0.05% concentration Make pin stream applications directly to the nests. To help avoid stings treat nests in the evening when the insects are inside the nest. For best results, check nests within one or two days to ensure complete kill. Remove and destroy treated nests to prevent emergence of newly hatched insects, secondary pest problems, and possible staining from honey. If removal is not possible, retreat nests if necessary.
BOXELDER BUGS	Apply directly to boxelder bugs that have congregated around doors and windows and similar areas where they may enter structures. Apply perimeter treatment as described above. For best results, treat infested trees early in the season. Tree spraying may require ornamental license - check with your local regulator.
CENTIPEDES, MILLIPEDES, PILLBUGS, SOWBUGS	Apply perimeter treatment to reduce the number of insects from entering the structure. Thoroughly treat mulched areas. Reduce moisture areas from around the structure.
FIRE ANTS	Use Rate: 0.05% concentration Thoroughly drench or inject the entire mound. When using injection treatment, you must treat the bottom of the mound. Apply dilution or foam using a downward and upward motion of the injector tool. For complete distribution, inject in several points throughout the mound. Treat new mounds as they appear. Do not use in pastures or areas of food crop production.

(continued)

FLEAS & TICKS	Make a perimeter treatment, concentrating on areas where pets rest or run. For ticks, treat likely harborage areas like along paths, tall grass and other vegetation. For best results, treat the entire area where ticks could be present. Begin treating in the spring. Ticks may be re-introduced from surrounding areas by host animals. Continue treating until frost to control both larvae and adult ticks.
FLIES	Use Rate: 0.025% to 0.05% concentration Apply low-volume mist or spray to outside surfaces of buildings, porches, patios, garages, fences, and other areas where flies rest. CLUSTER FLIES: Treat in late summer or fall, just before flies begin looking for winter shelter. Spray siding, eaves, around windows and doors, window screens, and attics. To help reduce infestations, treat cracks and crevices, attics and other indoor winter hiding places. Note: Cluster flies may appear during winter warm spells, while snow is still on the ground.
MOSQUITOES	Use Rate: 0.025% to 0.05% concentration (8 to 16 ml of TEMPO SC ULTRA per gal of water). Apply at the rate of one gal of dilution per 1,000 sq ft as a general spray around landscapes, lawns and buildings to control adult mosquitoes. For higher volume applications, TEMPO SC ULTRA may be diluted at lower concentrations and applied in greater volumes of water to deliver the desired amount of product per area (refer to Table 1: Mixing Directions for Outdoor/Perimeter Applications). Spray around landscape plants, turf and ground cover, under decks, around building foundations and other places where mosquitoes may rest.
SCORPIONS	Treat around piles of firewood and other areas where scorpions may hide. When possible, remove piles of materials that can harbor the pests. Apply a perimeter treatment.

STRUCTURAL PESTS				
WOOD-INFESTING PESTS				
<u>Beetles</u> Ambrosia Deathwatch	False powderpost Old house borer Powderpost	Carpenter ants Carpenter bees	<u>Termites</u> Subterranean Formosan	Drywood
WHERE TO APPLY	• Apply where pests have been seen or found, or can find shelter.			
USE RATE	• 0.05% -- 16 ml (0.54 fl oz) per gal water			
HOW TO APPLY (Also see Application section)	• Apply spray or foam as described in application section. • Avoid contact with treated surfaces until dry.			
WHEN TO APPLY	• When pests are seen or found • Re-apply every 7 to 10-days if needed.			
MIXING INFORMATION	• Add product when filling spray tank with water; shake or agitate mixture. • Diluted spray mixture can be stored overnight. Agitate before using. • Can be tank mixed with other pesticides currently registered for similar uses. To be sure of physical compatibility, first prepare a sample in a small jar using the proper proportions of water and products. Follow the most restrictive of labeling directions and precautions. Do not exceed label rates. • Foam applications: Mix end-use dilution with manufacturer's recommended volume of foaming agent for 0.025 to 0.05% concentration.			

APPLICATION FOR WOOD-INFESTING PESTS

CARPENTER ANTS	<u>In houses and other structures:</u> Apply around doors and windows and other places where carpenter ants may enter. Spray or foam into cracks & crevices, or spray, mist or foam through small drilled holes into voids where ants or nests are present. Use enough spray or foam to cover the area. Do not exceed 1 gal of dilution per 1,000 sq ft. Inside trees, utility poles, fencing and decking materials and similar structural members: Drill to find the interior infested cavity and inject dilution or foam using an appropriate treatment tool with a splashback guard. <u>Tunneling in soil:</u> Apply as a soil drench or inject the dilution or foam in the soil at intervals of 8 to 12 inches. Use 1 gal of dilution per 8 sq ft. Establish a uniform vertical barrier at the edge of walls, driveways or other hard surfaces where ants are tunneling beneath the surfaces. <u>Protecting firewood from carpenter ants:</u> Treat soil beneath firewood before stacking at 1 gal of dilution per 8 sq ft. Do not treat firewood with this product.
CARPENTER BEES	<u>In structures:</u> Spray or foam directly into gallery entrance holes. Leave entrance open for 24 hours to be sure that returning adult bees are killed. When there is no activity, the hole may be sealed with a suitable sealant.
WOOD-INFESTING BEETLES	On surfaces in homes, apartments, and other structures For small areas, brush or spray evenly on wood surfaces. For large or overhead areas, thoroughly cover wood surfaces with a coarse spray. Protect surfaces below the treated area from accidental dripping by covering with disposable plastic sheeting or other material that can be thrown away after treatment.

(continued)

	<u>In localized areas:</u> Apply dilution or foam to voids and galleries in damaged wood, in spaces between wooden structural members, and between the sill plate and foundation. Apply to inaccessible areas by drilling, and then injecting, spraying, or misting the dilution or foam into the damaged wood or wall voids.
DRYWOOD AND ABOVE-GROUND SUBTERRANEAN TERMITES	This type of application is intended to be a supplemental treatment for control of drywood and above-ground subterranean termites and other wood-infesting insects. <u>In localized areas:</u> Apply dilution or foam to voids and galleries in damaged wood, to spaces between wooden structural members, and between the sill plate and foundation. Apply to inaccessible areas by drilling, and then injecting, spraying, or misting the dilution or foam into damaged wood or wall voids. <u>Swarming or reproductive stages:</u> Apply dilution to the swarm or area on which the termites collect. Make applications to attics, crawl spaces, unfinished basements or man-made voids with a coarse fan spray to control exposed reproductive forms of termites. <u>Above-ground termites inside trees, utility poles, fencing and decking materials and similar structural members:</u> Drill to find the infested cavity and inject dilution or foam using an appropriate tool with a splashback guard. <u>Termite carton nests:</u> Inject dilution or foam into building voids or tree cavities using a pointed injection tool. Several injection points to varying depths may be needed. For best results, remove carton nest material when possible. <u>Protecting firewood from termites:</u> Treat soil beneath firewood before to stacking at 1 gal of dilution per 8 sq ft. Do not treat firewood with this product.

TURF PESTS

TEMPO SC ULTRA controls designated pests on landscape and recreational turf.

TURF PESTS			USE RATE
Ants Armyworms Crickets Cutworms	Earwigs Grasshoppers Japanese beetles (adult)	Sod webworms Ticks (including deer ticks)	4 to 8 ml (0.135 to 0.270 fl oz) per 1000 sq ft or 6 to 12 fl oz per acre Use high rate for severe infestations, faster knockdown, and longer residual.
Black turfgrass atenioides (adults) Bluegrass billbug (adult) Chinch bugs	Chiggers Fire ants (aids in control) Fleas	Hyperodes weevil (adult) Mole crickets Scorpions	8 ml (0.270 fl oz) per 1000 sq ft or 12 fl oz per acre
WHERE TO APPLY	For use on landscape and recreational turfgrass. Not for use on turf grown for research, commercial seed production, sale, or other commercial use. Not for use on golf course turf. Apply where pests have been seen or found, or can find shelter. Permitted areas include but are not limited to: home lawns, business complexes, apartment buildings, airports, cemeteries, parks, playgrounds, recreational fields, institutional grounds, athletic fields, and other landscape turf grass.		
HOW TO APPLY (Also see APPLICATION FOR TURF PESTS)	Use an accurately calibrated sprayer. Apply uniformly using spray nozzles with a coarse droplet size such as flood jet nozzles. Keep children and pets off treated areas until lawn has dried.		

(continued)

WHEN TO APPLY	Apply when pests first appear. Re-apply as needed based on pest reinfestation. Do not treat more than 6 times per year. Consult your State Cooperative Extension Office for the specific timing in your area.
MIXING INFORMATION	• Add product when filling spray tank with water; shake or agitate mixture. • Diluted spray mixture can be stored overnight. Agitate before using. Can be tank mixed with other pesticides currently registered for similar uses. Follow the most restrictive of labeling directions and precautions. Do not exceed label rates. Compatibility: Compatible with commonly used fungicides, miticides, liquid fertilizers and other insecticides. To be sure of physical compatibility, first prepare a sample in a small jar using the proper proportions of water and products.
RESTRICTIONS	• Do not apply this product by aerial application or by any type of chemigation system. • Do not apply when the average wind speed is greater than 15 mph. • Do not allow drift onto ponds, streams, or lakes. • Do not apply when turf grass areas are water logged or the soil is saturated with water (i.e., will not accept irrigation).

APPLICATION FOR TURF PESTS	
ARMYWORMS, CUTWORMS, SOD WEBWORMS	Do not water or mow for at least 24 hours following application.
BILLBUGS	Apply when adults first appear in the spring. In temperate climates, this application will also control overwintering chinch bugs.

(continued)

CHINCH BUGS	For best results, water the turf immediately after application to aid in the penetration of the active ingredient into the thatch layer where chinch bugs dwell.
TICKS (Including ticks that transmit Lyme Disease and Rocky Mountain Spotted fever)	For best results, treat the entire area where ticks could be present. Begin treating in the spring. Ticks may be re-introduced from surrounding areas by host animals. Continue treating until frost to control both larvae and adult ticks.

ORNAMENTAL PESTS

TEMPO SC ULTRA controls designated pests on trees, shrubs, foliage plants and flowers in outdoor landscaped areas and interior plantscapes where these plants are grown.

Ants	Cutworms	Leaf skeletonizers
Aphids	Elm leaf beetles	Mealybugs
Armyworms	Emerald Ash borer (Adult)*	Midges
Azalea caterpillars	Elm spanworms	Mosquitoes
Bagworms	Flea beetles	Oleander moth larvae
Black vine weevil (adult)	Flies	Orchid weevil
Boxelder bugs	Fungus gnats	Pear psylla
Bristly rose slugs	Grasshoppers	Peppertree psyllid
Budworms	Gypsy moth larvae	Pillbugs
California oakworms	Japanese beetles (adult)	Pine shoot moths
Cankerworms	June beetles (adult)	Pine tip moths
Casebearers	Lace bugs	Plant bugs
Clover mites	Leaf-feeding caterpillars	Redhumped caterpillars
Cockroaches	Leafhoppers	Sawfly larvae
Crickets	Leafrollers	Scale insects (crawler stages)

(continued)

Sowbugs	Tent caterpillars	Walnut caterpillars
Spiders	Thrips	Webworms
Spittlebugs	Ticks (including deer ticks)	Whiteflies
Striped beetles	Tussock moth larvae	Yellownecked caterpillars
Striped oakworms	Wasps	
*For emerald ash borer, make a minimum of two applications per season. The first application should be made in late May/early June or when adult beetles begin flying. Subsequent applications should be made approximately one month after the previous application.		
WHERE TO APPLY	Use in landscapes and interiorscapes. Permitted areas include but are not limited to: parks, recreational areas, athletic fields, institutional grounds, hotels, malls, office buildings, and airports. Not for use on plants or turf grown for research, commercial seed production, sale, or other commercial use. Not for use in greenhouses or nurseries. Apply where pests are seen, have been found, or can find shelter among trees, shrubs, evergreens, ground covers, flowers, and foliage plants. No phytotoxicity has been observed on a wide range of ornamentals tested under various environmental conditions. If in doubt, first try this product on a small sample.	
USE RATE	• 45 to 160 ml (1.5 to 5.4 ounces) TEMPO SC ULTRA per 100 gals of water • Use high rate for severe infestations, faster knockdown, and longer residual.	
HOW TO APPLY (Also see APPLICATION FOR ORNAMENTAL PESTS)	Apply as a dilute spray just to the point of run-off but do not allow excess run-off. For hard-to-wet foliage such as holly, pine, or ivy, adding a spreader-sticker may enhance knockdown and residual activity. For best results, spray thoroughly for good coverage including the underside of leaves for pests that feed or rest there.	

(continued)

	• Spray plants in bloom at times when pollinating insects are not present, such as early morning or late evening. • Remove any animal feeding dishes prior to treatment. Avoid contact with treated surfaces until dry.
WHEN TO APPLY	• Apply when pests first appear. Re-apply as needed based on pest infestation. • Consult your State Cooperative Extension Office for the specific timing in your area.
MIXING INFORMATION	• Add product when filling spray tank with water; shake or agitate mixture. • Diluted spray mixture can be stored overnight. Agitate before using. Can be tank mixed with other pesticides currently registered for similar uses. Follow the most restrictive of labeling directions and precautions. Do not exceed label rates. Compatibility: Compatible with commonly used fungicides, miticides, liquid fertilizers and other insecticides. To be sure of physical compatibility, first prepare a sample in a small jar using the proper proportions of water and products.
RESTRICTIONS	• Do not apply to any food crop. • Do not allow drift onto ponds, streams or lakes.

APPLICATION FOR ORNAMENTAL PESTS

ANTS, CRICKETS, SPIDERS, MIDGES, WASPS AND FLIES	Apply as a general spray, concentrating on damp areas beneath shrubbery, around foundations and in areas of tall grass or weeds in ornamental areas and areas adjacent to buildings.
BAGWORMS	Spray when bagworms first begin to hatch. If possible, apply directly to the larvae. Young larvae are easier to control.

(continued)

BLACK VINE WEEVIL (adult)	Use rate: 60 to 160 ml (2 to 5.4 fl oz) per 100 gals Apply as needed when adults and damage are first noticed in spring and at 3 to 4 week intervals while weevils and notching are observed. Addition of a spreader/sticker at recommended rates may enhance control of insects on certain species of ornamentals having difficult to wet foliage.
COCKROACHES	Spray hiding or resting places such as under shrubbery, in ground covers and mulching materials, in hollow trees, etc.
SCALE (CRAWLERS)	Spray thoroughly, including foliage, trunk, limbs, and twigs.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage:	• Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, or feed. • Protect from freezing. • Store in original containers and out of the reach of children, preferably in a locked storage area. • Handle and open container in a manner as to prevent spillage.
Pesticide Disposal:	• Pesticide wastes are hazardous. • Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. • If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

(continued)

Container Disposal:	Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Securely wrap container in several layers of newspaper and discard in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.
Spills:	If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. • Do not walk through spilled material. • Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides above. • Keep unauthorized people away. • You may contact the Bayer Environmental Science Emergency Response Team for decontamination procedures or any other assistance at 1-800-334-7577.

APPENDIX

Use TEMPO SC ULTRA as a general surface, spot, mist, or crack and crevice treatment. Permitted areas of use in buildings, structures, and modes of transport include, but are not limited to:

- | | | |
|---|---------------------------------|--|
| • apartment buildings | • granaries | • nursing homes |
| • bakeries | • greenhouses (structures only) | • pickle factories |
| • bottling facilities | (non-commercial) | • poultry houses |
| • breweries | • hospitals | • processing and packing plants |
| • buses | • hotels | • restaurants |
| • cafeterias | • houses | • schools |
| • candy plants | • industrial buildings | • spice plants |
| • cereal processing and manufacturing plants | • kitchens | • stores |
| • correctional facilities | • laboratories | • warehouses |
| • dairies and dairy product processing plants | • livestock housing | • wineries and similar structures |
| • frozen food plants | • pet kennels | • transportation vessels such as aircraft - cargo areas only, buses, trucks, trailers, rail cars, and marine vessels |
| • grain mills | • manufacturing establishments | |
| | • mausoleums | |
| | • meat and produce canneries | |

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product.

If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience LP. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSOURCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSOURCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSOURCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

Bayer (reg'd), the Bayer Cross (reg'd), Tempo® and Backed by Bayer™ are trademarks of Bayer.



Produced for:

Bayer Environmental Science

A Division of Bayer CropScience LP

2 T. W. Alexander Drive

Research Triangle Park, NC 27709

2.5"

3"

TEMPO[®] SC

ULTRA

Insecticide

**For Pest Management
Professionals and Commercial
Use Only.**

ACTIVE INGREDIENT:

β-Cyfluthrin, Cyano(4-fluoro-
3-phenoxyphenyl) methyl
-(2,2-dichloroethenyl)
-2,2-dimethyl-
cyclopropanecarboxylate . . . 11.8%

OTHER INGREDIENTS: . . . 88.2%

100.0%

Contains 1 lb
cyano(4-fluoro-3-phenoxyphenyl)
methyl 3-(2,2-
dichloroethenyl)-2,2-dimethylcyclo
propanecarboxylate per gallon.

SHAKE WELL BEFORE USING

EPA Reg. No. 432-1363

STOP - Read The Label Before Use.

Keep out of reach of children.

CAUTION

Net Contents:

240 mL (8.12 fl oz)

4031958

3549856C 100504AV1